

Threatened Species of the Northern Territory

SANDHILL DUNNART *Sminthopsis psammophila*

Conservation status

Australia: Endangered.
Northern Territory: Data Deficient.



Sandhill dunnart. (Photo: P. Canty NPWSA)

Description

The sandhill dunnart is a medium sized carnivorous marsupial with a body mass of 26-40 g. The upperparts are a plain grey to buff with darker hairs interspersed throughout. There is a dark triangle of fur on the crown and forehead. The face and flanks are buff, the eyes are dark. The underparts and feet are whitish. The tail length is longer than the head-body length. The tail is grey above and darker below with a crest of short, black hairs near the tip.


Distribution

The first specimen of the sandhill dunnart was collected by members of the Horn Expedition in 1894 near Lake Amadeus between Kurtitina Well and Uluru (Parker 1973). Since that time it has not been seen in the Territory, although remains of the species have been identified in owl pellets collected from Uluru and Kata Tjuta (Baynes and Johnson 1996).

Outside of the Northern Territory, the species has been recorded on the Eyre Peninsula (South Australia) and in the Great Victoria Desert (South Australia and Western Australia). (Pearson and Robinson 1990). A total of 60 individuals have been

captured at all sites up to 2000 (Churchill 2001).

Conservation reserves where reported:
None.



Known locations of the long-tailed dunnart.
o = pre 1970; • = post 1970

Ecology

The Northern Territory specimen came from sand dune country in an area covered by spinifex hummocks with groves of desert oaks in the swales. On the Eyre Peninsula, the sandhill dunnart occupies sand ridges covered by hummock grassland and mallee-broombush shrub. In the Great Victoria

Desert and at Yellabinna it occurs in sandy environments, either sand plains or dunes. The vegetation occupied consists of low woodland or low open woodland with a diverse shrub understorey and a ground cover of at least 20% spinifex hummocks (Pearson and Robinson 1990). The species is insectivorous and breeding takes place in spring and early summer (Pearson 1995).

Conservation assessment

The sandhill dunnart has been recorded only once in the Northern Territory and that was over 100 years ago. However, the small amount of survey work carried out in the vicinity of Lake Amadeus where the specimen was obtained means that it is premature to regard it as regionally extinct. Therefore, the status of the sandhill dunnart in the Northern Territory is **Data deficient**.

Threatening processes

Inappropriate fire regimes and clearance of land for agriculture are major threats faced by the sandhill dunnart in southern Australia (Churchill 2001). In particular, the species prefers large, mature spinifex clumps in which it builds a small nest chamber composed of spinifex needles. Spinifex reaches the size and structure preferred by sandhill dunnarts 5-10 years after fire and this type of spinifex constitutes only 5% of the available spinifex hummocks on sites occupied by the dunnart on the Eyre Peninsula (Churchill 2001). Therefore, fire regimes that threaten the preferred nest sites of the sandhill dunnart may represent a long term threat to the species. Predation by feral cats and foxes may also be a threat.

Conservation objectives and management

There is a national Recovery Plan for the sandhill dunnart (Churchill 2001), but no existing management program in the Northern Territory.

Any future survey of the Great Sandy Desert bioregion, particularly in the vicinity of Lake Amadeus, should include targeted surveys for this species.

This species is likely to benefit from fire management that maintains an adequate proportion of appropriately-aged spinifex.

Compiled by

Chris Pavey
[May 2006]

References

- Baynes, A., and Johnson, K.A. (1996). The contributions of the Horn Expedition and cave deposits to knowledge of the original mammal fauna of central Australia. In *Exploring Central Australia: Society, the Environment and the 1894 Horn Expedition* (eds S.R. Morton and D.J. Mulvaney.) pp. 168-186. (Surrey Beatty and Sons, Sydney.)
- Churchill, S. (2001). *Recovery plan for the sandhill dunnart (Sminthopsis psammophila)*. (Environment Australia, Canberra.)
- Parker, S.A. (1973). An annotated checklist of the native land mammals of the Northern Territory. *Records of the South Australian Museum* **16**, 1-57.
- Pearson, D.J. (1995). Sandhill dunnart. In *The Mammals of Australia*. 2nd Edition (ed. R. Strahan.) pp. 154-155. (Australian Museum, Sydney.)
- Pearson, D.J., and Robinson, A.C. (1990). New records of the sandhill dunnart, *Sminthopsis psammophila* (Marsupialia: Dasyuridae) in South and Western Australia. *Australian Mammalogy* **13**, 57-59.